

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Figure 5

Use of A2/Melan-A peptide tetramers for monitoring of antigen peptide-driven expansion of Melan-A specific CTL from precursors in the peripheral blood lymphocytes of a melanoma patient.

Stimulating peptide :



→ α CD8



Figure 6

Assay of lytic activity of PBMC from a melanoma patient after three rounds of in vitro stimulation with Melan-A derived peptides.





Figure 7

Melan-A specific lytic activity of fluorescently sorted tetramer⁺ lymphocytes

stimulated with:



lymphocytes to target cell ratio